

QUASI-ISOMETRIC EMBEDDINGS INTO DIFFEOMORPHISM GROUPS

MICHAEL BRANDENBURSKY AND JAREK KĘDRA

ABSTRACT. Let M be a smooth compact connected oriented manifold of dimension at least two endowed with a volume form. Assuming certain conditions on the fundamental group $\pi_1(M)$ we construct quasi-isometric embeddings of either free Abelian or direct products of non-Abelian free groups into the group of volume preserving diffeomorphisms of M equipped with the L^p metric induced by a Riemannian metric on M .

1. INTRODUCTION

1.A. The L^p -metric. Let M be a compact connected and oriented Riemannian manifold and let $\text{Diff}(M, \mu)$ denote the group of smooth diffeomorphisms of M acting by the identity on a neighborhood of the boundary and preserving the volume form μ induced by the metric. Unless otherwise stated we assume that $\text{Diff}(M, \mu)$ is equipped with the Whitney C^∞ -topology.

In the present paper we study the geometry of the identity component $\text{Diff}_0(M, \mu)$ of the above group endowed with the right invariant L^p -metric. It is defined as follows. Let

$$\mathcal{L}_p\{g_t\} := \int_0^1 dt \left(\int_M |\dot{g}_t(x)|^p \mu \right)^{\frac{1}{p}}$$

be the L^p -length of a smooth isotopy $\{g_t\}_{t \in [0,1]} \subset \text{Diff}_0(M, \mu)$, where $|\dot{g}_t(x)|$ denotes the length of the tangent vector $\dot{g}_t(x) \in T_x M$ induced by the Riemannian metric. Observe that this length is right-invariant, that is, $\mathcal{L}_p\{g_t \circ f\} = \mathcal{L}_p\{g_t\}$ for any $f \in \text{Diff}(M, \mu)$. It defines a right-invariant metric on $\text{Diff}_0(M, \mu)$ by

$$\mathbf{d}_p(g_0, g_1) := \inf_{g_t} \mathcal{L}_p\{g_t\},$$

where the infimum is taken over all paths from g_0 to g_1 .

If $p = 2$ then the group $\text{Diff}_0(M, \mu)$ is in fact equipped with a Riemannian metric inducing the above L^2 -length. The geodesics of this metric are the solutions of the equations of the flow of an incompressible fluid [1], which makes the $p = 2$ case the most interesting. It is known that if M is a simply connected Riemannian manifold of dimension at least

three then the L^2 -diameter of the group $\text{Diff}_0(M, \mu)$ is finite [18]. On the other hand Eliashberg and Ratiu [10] proved that this diameter is infinite for surfaces and for manifolds with positive first Betti number. See Arnol'd-Khesin [2] and Khesin-Wendt [14, Section 3.6] for a detailed discussion.

1.B. The main result. A map $\psi: (X_1, d_1) \rightarrow (X_2, d_2)$ between metric spaces is called *large scale Lipschitz* [17, Remark 1.9] if there exist constants $A, B \geq 0$ such that

$$d_2(\psi(x), \psi(y)) \leq A \cdot d_1(x, y) + B.$$

Let $m \in M$ be a reference point. Let $\text{ev}_m: \text{Diff}_0(M, \mu) \rightarrow M$ be the evaluation map defined by $\text{ev}_m(f) := f(m)$ and let $G_\mu \subset \pi_1(M)$ be the image of the homomorphism induced by ev_m . It is easy to prove that G_μ is contained in the center of $\pi_1(M)$. The subgroup G_μ is called the *Gottlieb group* associated with the volume form μ because the groups of similar origin were first studied by Gottlieb in [11].

Let $\text{Diff}(M, \mu, m) \subset \text{Diff}_0(M, \mu)$ be the isotropy of the reference point m . Let us define a homomorphism

$$\Phi: \text{Diff}(M, \mu, m) \rightarrow \pi_1(M)/G_\mu$$

as follows. Let $g \in \text{Diff}(M, \mu, m)$ and let $\{g_t\} \subset \text{Diff}_0(M, \mu)$ be a smooth isotopy from the identity to g . The value $\Phi(g)$ is represented by the loop $\{g_t(m)\}$ and it is straightforward to show that Φ is well defined.

Let $B(m, r) \subset M$ be a ball of radius $r > 0$ centered at a reference point $m \in M$. Let $\text{Diff}(M, \mu, B(m, r))$ denote the subgroup of $\text{Diff}_0(M, \mu)$ consisting of diffeomorphisms preserving the ball $B(m, r)$ pointwise. The metric on the group $\text{Diff}(M, \mu, B(m, r))$ is induced from the L^p -metric on $\text{Diff}_0(M, \mu)$. The following is our main technical result which is proven in Section 2.

Theorem 1.1. *Let M be a compact connected and oriented Riemannian manifold. For all small enough $r > 0$ the homomorphism*

$$\Phi: \text{Diff}(M, \mu, B(m, r)) \rightarrow \pi_1(M)/G_\mu$$

is surjective and large scale Lipschitz with respect to the L^p -metric on the group $\text{Diff}(M, \mu, B(m, r))$ and the word metric on $\pi_1(M)/G_\mu$.

Recall that the word norm on a group Γ generated by a symmetric finite set $S \subset \Gamma$ is defined by

$$|\gamma|_S := \min\{k \in \mathbb{N} \mid \gamma = s_1 \dots s_k \text{ where } s_i \in S\}.$$

The word metric is defined by $d_S(\gamma_1, \gamma_2) := |\gamma_1(\gamma_2)^{-1}|_S$. It is right-invariant and it depends on the choice of a finite generating set up to a quasi-isometry (defined below) [5, Example 8.17].

1.C. Applications. Let (X_1, d_1) and (X_2, d_2) be two metric spaces. A function $f: X_1 \rightarrow X_2$ is a *quasi-isometric embedding* if there exist two constants $A \geq 1$ and $B \geq 0$ such that

$$(1) \quad \frac{1}{A} d_1(x, y) - B \leq d_2(f(x), f(y)) \leq A d_1(x, y) + B.$$

In the case when (X_1, d_1) and (X_2, d_2) are metric groups, we require f to be an *injective homomorphism*. We say that f is a *quasi-isometry* if, in addition to (1), there exists a constant $C \geq 0$ such that for every $u \in X_2$ there exists $x \in X_1$ with the property

$$d_2(u, f(x)) \leq C.$$

Let Γ be a group equipped with the word metric associated with a finite generating set $S \subset \Gamma$. An element γ is called *undistorted* in Γ if there exists a positive constant $C > 0$ such that

$$|\gamma^n|_S \geq C \cdot n.$$

Otherwise, γ is called *distorted*. These properties do not depend on the choice of a finite generating set.

Theorem 1.2. *Let M be a compact connected and oriented Riemannian manifold of dimension at least two. If $\pi_1(M)/G_\mu$ contains an undistorted element then $(\text{Diff}_0(M, \mu), \mathbf{d}_p)$ contains quasi-isometrically embedded free Abelian group of an arbitrary rank. In particular, the metric group $(\text{Diff}_0(M, \mu), \mathbf{d}_p)$ has infinite diameter.*

This result generalizes a theorem of Eliashberg and Ratiu [10] where they prove the infiniteness of the diameter under the assumption that the first Betti number of M is positive. Notice that there exist compact oriented manifolds with the fundamental group isomorphic to an arbitrary finitely presented group. Such a group can be chosen to have finite abelianization (hence the first Betti number is equal to zero) and usually groups have undistorted elements. For example, $\text{SL}(2, \mathbf{Z})$ has finite abelianization. Moreover, if $g \in \text{SL}(2, \mathbf{Z})$ has an eigenvalue λ such that $|\lambda| \neq 1$ then the cyclic group generated by g is undistorted (as everybody knows).

In fact, no finitely presented group with all its elements distorted is known. On the other hand, Osin [16] constructs an example of an infinite finitely generated group with exactly two conjugacy classes. In particular, such a group does not have undistorted elements.

Theorem 1.3. *Let M be a compact connected and oriented Riemannian manifold of dimension at least three. If $\pi_1(M)/G_\mu$ contains quasi-isometrically embedded non-Abelian free group then $(\text{Diff}_0(M, \mu), \mathbf{d}_p)$ contains quasi-isometrically embedded direct product of any finite number of free groups of arbitrary ranks.*

Essentially, the idea of the proof is to appropriately embed disjoint copies of the figure eight into M and suitably apply Theorem 1.1. Thus the situation is a bit different for surfaces, where we get the following slightly weaker statement.

Theorem 1.4. *Let $\Sigma_{g,k}$ be a compact connected and oriented surface of genus g with k boundary components. Then $\text{Diff}_0((\Sigma_{g,k}, \mu), \mathbf{d}_p)$ contains quasi-isometrically embedded direct product of $2g+k-2$ copies of finitely generated non-Abelian free groups of arbitrary ranks.*

Remark 1.5. Let \mathbb{D}^2 be a unit Euclidean disc in \mathbf{R}^2 , i.e. in our notation \mathbb{D}^2 is diffeomorphic to $\Sigma_{0,1}$. In [3] Benaim and Gambaudo showed, using a different method, that the group $\text{Diff}((\mathbb{D}^2, \mu), \mathbf{d}_p)$ contains quasi-isometrically embedded finitely generated free or free Abelian group of arbitrary rank. Crisp and Wiest proved the same fact for planar right-angled Artin groups [7]. We also would like to mention that M. Kapovich proved that any right-angled Artin group embeds into the group of Hamiltonian diffeomorphisms of any symplectic manifold (M, ω) , see [12]. However, it is not known whether the embedding he constructs in [12] is quasi-isometric with respect to L^p -metric.

1.D. Examples.

Example 1.6. Since the Artin pure braid group \mathbf{P}_3 on three strands is isomorphic to $\mathbf{F}_2 \times \mathbf{Z}$ (see the proof of Theorem 1.16 in [13]) it embeds quasi-isometrically into $\text{Diff}_0(M, \mu)$ where M is as in Theorem 1.3.

Example 1.7. If a subgroup Γ of a direct product of n free groups has finitely generated homology up to degree n then Γ contains a finite index subgroup isomorphic to a direct product of at most n free groups [6]. On the other hand, there are examples of finitely presented subgroups of a direct product of free groups which have more complicated finiteness properties. More concretely, the kernel Γ of the homomorphism $(\mathbf{F}_2)^n \rightarrow \mathbf{Z}$ sending each generator to one is of type FP_n but not FP_{n+1} . These examples are known as *Bieri-Stallings groups*, see Bestvina-Brady [4, Example 6.3].

Suppose that $\Gamma \rightarrow \mathbf{F}_2 \times \dots \times \mathbf{F}_2$ is the inclusion of a finitely presented Bieri-Stallings group. It follows from the proof of Theorem 11.7 of Dison's thesis [9] that this inclusion is a quasi-isometric embedding. Indeed, since the quotient is isomorphic to \mathbf{Z} , its isoperimetric function is linear and hence, by Dison's Lemma 9.5 in [9] the distortion function of the above inclusion is linear.

Recall that if $\psi: H \rightarrow G$ is an injective homomorphism of metric groups then its distortion function $\Delta: \mathbf{R}_+ \rightarrow \mathbf{R}_+$ is defined by

$$\Delta(r) := \max\{d_H(1, h) \mid d_G(1, \psi(h)) < r\}.$$

Observe that the distortion function is linear if and only if ψ is a quasi-isometric embedding.

Example 1.8. Let $M \subset \mathbf{R}^3$ be a closed domain with free non-Abelian fundamental group. The geometry of $(\text{Diff}_0(M, \mu), \mathbf{d}_p)$ models the behavior of an incompressible fluid filling a tank of the shape of M . Theorem 1.3 describes a large scale complexity of mixing such a fluid.

1.E. The symplectic case. If (M, ω) is a symplectic manifold, then the group $\text{Diff}_0(M, \mu)$ in all the results above can be replaced either by the group $\text{Symp}_0(M, \omega)$ of symplectic diffeomorphisms isotopic to the identity, or by the group $\text{Ham}(M, \omega)$ of Hamiltonian diffeomorphisms, see Remark 2.2 in the proof of Theorem 1.1.

2. PROOFS

2.A. An abstract lemma. Let (G, d_G) be a metric group with the identity element 1_G . For $g \in G$ we set

$$\|g\|_G := d_G(1_G, g) \quad \text{and} \quad \text{diam}(G) := \sup_{g \in G} \|g\|_G.$$

Lemma 2.1. *Let (G, d_G) , (H, d_H) , (K, d_K) be three metric groups, such that H is finitely generated and d_H is a word metric w.r.t. some finite generating set S . Suppose that $\Phi: G \rightarrow K$ is a large scale Lipschitz homomorphism. Let $\Psi: H \rightarrow G$ be a homomorphism, such that $\Phi\Psi: H \rightarrow K$ is a quasi-isometric embedding. Then Ψ is a quasi-isometric embedding.*

Proof. The homomorphism Ψ is injective, because $\Phi\Psi$ is injective. Let $h \in H$. The homomorphism $\Phi\Psi$ is a quasi-isometric embedding, hence there exist two constants $A_1 \geq 0$ and $B_1 \geq 0$ such that

$$(2) \quad A_1 \|h\|_H - B_1 \leq \|\Phi\Psi(h)\|_K$$

The homomorphism Φ is a large scale Lipschitz, hence there exist two constants $A_2 > 0$ and $B_2 > 0$ such that

$$(3) \quad \|\Phi\Psi(h)\|_K \leq A_2 \|\Psi(h)\|_G + B_2.$$

Combining inequalities (2) and (3) we get

$$(4) \quad \frac{A_1}{A_2} \|h\|_H - \left(\frac{B_1 + B_2}{A_2} \right) \leq \|\Psi(h)\|_G.$$

Let k be such that $\|h\|_H = k$. The group H is finitely generated, hence $h = h_1 \cdot \dots \cdot h_k$, where $h_i \in S$ for each $1 \leq i \leq k$. Denote by

$$M_\Psi := \max\{\|\Psi(h_1)\|_G, \dots, \|\Psi(h_k)\|_G\}.$$

It follows that

(5)

$$\|\Psi(h)\|_G = \|\Psi(h_1) \cdot \dots \cdot \Psi(h_k)\|_G \leq \sum_{i=1}^k \|\Psi(h_i)\|_G \leq M_\Psi \cdot k = M_\Psi \|h\|_H.$$

Inequalities (4) and (5) conclude the proof of the lemma. \square

2.B. Proof of Theorem 1.1. In the first part we prove the surjectivity of the homomorphism $\Phi: \text{Diff}(M, \mu, B(m, r)) \rightarrow \pi_1(M)/G_\mu$.

Let $S' := \{[\gamma_1], \dots, [\gamma_k]\} \subset \pi_1(M)$ be a symmetric generating set of the fundamental group of M such that each representative γ_i is a simple closed curve. It follows from the tubular neighborhood theorem that for each γ_i there exists $r_i > 0$, the standard $(n-1)$ -dimensional ball $B_{r_i}^{n-1} \subset \mathbf{R}^n$ of radius $r_i > 0$, and a volume-preserving embedding

$$\text{emb}_i : B_{r_i}^{n-1} \times \mathbf{S}^1 \hookrightarrow M,$$

such that $\text{emb}_i|_{\{0\} \times \mathbf{S}^1}$ is the curve γ_i and $r_i = d_M(\gamma_i, \text{emb}_i|_{\partial B_{r_i}^{n-1} \times \mathbf{S}^1})$. The volume form on the product $B_{r_i}^{n-1} \times \mathbf{S}^1$ is the standard Euclidean volume and d_M denotes the distance on M induced by the Riemannian metric.

Let $r = \min_{1 \leq i \leq k} r_i$. Then $\text{emb}_i : B_r^{n-1} \times \mathbf{S}^1 \hookrightarrow M$ is volume-preserving for each i and $\text{emb}_i|_{\{0\} \times \mathbf{S}^1} = \gamma_i$. It is straightforward to construct a smooth isotopy of volume-preserving diffeomorphisms

$$g_t : B_r^{n-1} \times \mathbf{S}^1 \rightarrow B_r^{n-1} \times \mathbf{S}^1$$

between $g_0 = \text{Id}$ and g_1 such that:

- For each $t \in [0, 1]$ the diffeomorphism g_t equals to the identity in the neighborhood of $\partial B_r^{n-1} \times \mathbf{S}^1$, and the time-one map g_1 is equal to the identity on $B_{r'}^{n-1} \times \mathbf{S}^1$, where $0 < r' < r$.
- Each diffeomorphism g_t preserves the foliation of $B_r^{n-1} \times \mathbf{S}^1$ by the circles $\{x\} \times \mathbf{S}^1$ and for every $x \in B_{r'}^{n-1}$ the restriction $g_t : \{x\} \times \mathbf{S}^1 \rightarrow \{x\} \times \mathbf{S}^1$ is the rotation by $2\pi t$. It follows that the time-one map g_1 equals the identity on $B_{r'}^{n-1} \times \mathbf{S}^1$.

We identify $B_r^{n-1} \times \mathbf{S}^1$ with its image with respect to the embedding emb_i . Then we extend each isotopy g_t by the identity on $M \setminus B_r^{n-1} \times \mathbf{S}^1$ obtaining smooth isotopies $g_{t,i} \in \text{Diff}_0(M, \mu)$. This shows that every

representative γ_i of a generator of the fundamental group of M arises as simple closed curve $\{g_{t,i}(m)\}$ and hence the homomorphism

$$\Phi: \text{Diff}(M, \mu, B(m, r)) \rightarrow \pi_1(M)/G_\mu$$

is surjective.

Remark 2.2. Notice that if M is a symplectic manifold then the above isotopies can be constructed to be Hamiltonian.

Let $\Pi: \pi_1(M) \rightarrow \pi_1(M)/G_\mu$ be the projection homomorphism. Consequently $S := \Pi(S')$ is a finite generating set for the quotient $\pi_1(M)/G_\mu$. Let $\Pi_M: M_\bullet \rightarrow M$ be the Riemannian covering associated with Π . This means that the metric on M_\bullet is induced from the Riemannian metric on M . The corresponding distance will be denoted by d_\bullet .

Now we shall prove that Φ is a large scale Lipschitz map. That is, we show that there exist positive constants A and B independent of g such that

$$A \cdot \|g\|_p + B \geq \|\Phi(g)\|_S,$$

where $\|g\|_p := \mathbf{d}_p(g, \text{Id})$ is the L^p -norm of the diffeomorphism g .

Let $g \in \text{Diff}(M, \mu, B(m, r))$ and let $\{g_t\}_{t \in [0,1]} \in \text{Diff}_0(M, \mu)$ be an isotopy from the identity to g . It follows from the Hölder inequality that $\|g\|_p \geq C_p \cdot \|g\|_1$, where C_p is some positive constant independent of g . Hence it is enough to prove the statement for $p = 1$.

Let $m_\bullet \in \Pi_M^{-1}(m)$, and let $\{g_{\bullet,t}(m_\bullet)\}$ be the lift of $\{g_t(m)\}$ starting at the point m_\bullet . The manifold M is compact, hence by the Švarc-Milnor lemma [5, 15], the inclusion of the orbit of m_\bullet with respect to the deck transformation group $\pi_1(M)/G_\mu$ defines a quasi-isometry

$$\pi_1(M)/G_\mu \stackrel{q.i.}{\cong} (M_\bullet, d_\bullet).$$

In particular, it means that there exist positive constants A', B' , such that

$$(6) \quad d_\bullet(m_\bullet, g_{\bullet,1}(m_\bullet)) \geq A' \|\Phi(g)\|_S - B'.$$

Let $x \in B(m, r) \subset M$. We claim that the length of the flow-line $g_t(x)$ is bounded by the distance $d_\bullet(m_\bullet, g_{\bullet,1}(m_\bullet))$ up to the diameter of the ball. To see this, consider the lift of each flow-line $g_t(x)$ starting at a ball of radius r in M_\bullet centered at m_\bullet and observe that such a lift ends in a ball of radius r centered at $g_{\bullet,1}(m_\bullet)$. Indeed, let $\alpha: [0, 1] \rightarrow B(m, r)$ be a path between x and m . Then the map

$$H: [0, 1] \times [0, 1] \rightarrow M,$$

defined by $H(t, s) = g_t(\alpha(s))$ is a homotopy from $\{g_t(m)\}$ to $\{g_t(x)\}$. Lifting this homotopy shows that the lift of $\{g_t(x)\}$ finishes at the ball

$B(g_{\bullet,1}(m_\bullet), r)$. Finally, we obtain that

$$(7) \quad \text{Length}(g_t) := \int_0^1 |\dot{g}_t(x)| dt \geq d_\bullet(m_\bullet, g_{\bullet,1}(m_\bullet)) - 2r.$$

as claimed. Combining inequalities (6) and (7) we get that

$$\text{Length}(g_t(x)) \geq A' \|\Phi(g)\|_S - (B' + 2r)$$

for every $x \in B(m, r)$. Hence by Fubini theorem and the above inequality we have

$$\begin{aligned} \mathcal{L}_1(\{g_t\}) &= \int_0^1 dt \left(\int_M |\dot{g}_t(x)| d\mu \right) \\ &= \int_M \mu \left(\int_0^1 |\dot{g}_t(x)| dt \right) \\ &\geq \text{vol}(B(m, r)) \cdot \min_{x \in B(m, r)} \text{Length}(g_t(x)) \\ &\geq \text{vol}(B(m, r)) \cdot A' \|\Phi(g)\|_S - \text{vol}(B(m, r)) \cdot (B' + 2r). \end{aligned}$$

Since the above inequalities hold for any isotopy $\{g_t\}_{t \in [0,1]}$ between the identity and g , we obtain that

$$\|\Phi(g)\|_S \leq A \cdot \|g\|_p + B,$$

where $A = (C_p \cdot \text{vol}(B(m, r))^{-1} \cdot A'$ and $B = \frac{B' + 2r}{A}$ and this concludes the proof. \square

2.C. Proof of Theorem 1.2. Let $n \in \mathbf{N}$ be a positive integer. Recall that we need to prove that there exists a quasi-isometric embedding of a free Abelian group of rank n into $\text{Diff}_0(M, \mu)$.

Assume first that the dimension of M is at least three. Let m_1, \dots, m_n be distinct points in the interior of M and let $r > 0$ be such that the balls $B(m_i, r)$ of radius r centered at m_i are pairwise disjoint. Let $\gamma_{i,j}$ be simple closed curves representing the generators of $\pi_1(M, m_i)$. We also assume that γ_{i_1, j_1} is disjoint from γ_{i_2, j_2} whenever $i_1 \neq i_2$. We choose r small enough such that the tubular neighborhood of radius r of the above generators are disjoint.

Let $G_i \subset \pi_1(M, m_i)$ be the corresponding Gottlieb group. The groups $\pi_1(M, m_i)/G_i$ are pairwise isomorphic. Let $\gamma_i \in \pi_1(M, m_i)/G_i$ be an undistorted element which exists according to the hypothesis. Let

$$h: \mathbf{Z}^n \rightarrow \pi_1(M, m_1)/G_1 \times \dots \times \pi_1(M, m_n)/G_n$$

be a homomorphism defined by

$$h(k_1, \dots, k_n) := (\gamma_1^{k_1}, \dots, \gamma_n^{k_n}).$$

It immediately follows from the fact that each γ_i is undistorted that h is a quasi-isometric embedding. Let $\text{Diff}(M, \mu, \sqcup_i B(m_i, r)) \subset \text{Diff}_0(M, \mu)$

be the subgroup consisting of diffeomorphisms preserving the disjoint union of balls $B(m_i, r)$ pointwise. Let

$$\Phi_i: \text{Diff}(M, \mu, B(m_i, r)) \rightarrow \pi_1(M, m_i)$$

be the homomorphism defined in Theorem 1.1. Consider a homomorphism

$$\tilde{\Phi}: \text{Diff}(M, \mu, \sqcup_i B(m_i, r)) \rightarrow \pi_1(M, m_1)/G_1 \times \dots \times \pi_1(M, m_n)/G_n$$

which is the composition of the (diagonal) inclusion

$$\iota: \text{Diff}(M, \mu, \sqcup_i B(m_i, r)) \hookrightarrow \prod_i \text{Diff}(M, \mu, B(m_i, r))$$

followed by the product homomorphism

$$\prod_i \Phi_i: \prod_i \text{Diff}(M, \mu, B(m_i, r)) \rightarrow \prod_i \pi_1(M, m_i)/G_i.$$

Since the inclusion ι is an isometric embedding and the $\prod_i \Phi_i$ is large scale Lipschitz, according to Theorem 1.1, we obtain that $\tilde{\Phi}$ is a large scale Lipschitz homomorphism.

Let $g_i \in \text{Diff}(M, \mu, B(m_i, r))$ be an element such that $\Phi_i(g_i) = \gamma_i$ and g_i is supported in the union of the tubular neighborhoods of the loops representing the generators of $\pi_1(M, m_i)$ constructed in the beginning of the proof. It follows that the supports of g_i and g_j are disjoint if $i \neq j$. The existence of g_i follows from the proof of Theorem 1.1. Let

$$\Psi: \mathbf{Z}^n \rightarrow \text{Diff}(M, \mu, \sqcup_i B(m_i, r)) \subset \text{Diff}_0(M, \mu)$$

be defined by

$$\Psi(k_1, \dots, k_n) := g_1^{k_1} \circ \dots \circ g_n^{k_n}.$$

It is well defined because g_i have pair-wise disjoint supports. Recall that we have that

$$h = \tilde{\Phi} \circ \Psi: \mathbf{Z}^n \rightarrow \prod_i \pi_1(M, m_i)/G_i$$

and we know that h is a quasi-isometric embedding and $\tilde{\Phi}$ is large scale Lipschitz. Consequently the map Ψ is a quasi-isometric embedding according to Lemma 2.1.

Let us now consider the two-dimensional case. Let $M = \Sigma_{g,k}$ be a compact oriented surface of genus g with k boundary components. Observe that $\pi_1(\Sigma_{g,k})/G_\mu$ is trivial if either $g = 0$ and $k \leq 2$ or $g = 1$ and $k = 0$. Otherwise it is either free non-Abelian group or the fundamental group of a closed oriented surface. In each case it is straightforward to define an embedding

$$\text{emb}: \mathbf{S}^1 \times [0, 2n] \rightarrow M$$

such that each loop $\text{emb}(\mathbf{S}^1 \times \{t\})$ represents an undistorted element in $\pi_1(M, \text{emb}(1, t))$. Let $g_i: M \rightarrow M$, for $i = 1, \dots, n$ be an area preserving diffeomorphism satisfying each of the following conditions:

- it is supported in $\text{emb}(\mathbf{S}^1 \times (2i - 2, 2i))$;
- it preserves the ball $\text{emb}(B_i)$, where $B_i \subset \mathbf{S}^1 \times [0, 2n]$ is a ball of diameter one centered at $(1, 2i - 1)$;
- it is the time one map of an isotopy from the identity which acts as the full rotation on the loop $\text{emb}(\mathbf{S}^1 \times \{2i - 1\})$.

As in the previous part the homomorphism $\Psi: \mathbf{Z}^n \rightarrow \text{Diff}_0(M, \mu)$ defined by $\Psi(k_1, \dots, k_n) := g_1^{k_1} \circ \dots \circ g_n^{k_n}$ is the required quasi-isometric embedding. \square

2.D. Proof of Theorem 1.3. This proof is a modification of the proof of Theorem 1.2 for three dimensional M where the cyclic group \mathbf{Z} is replaced by a non-Abelian free group \mathbf{F}_2 on two generators. More precisely, let $f_i, g_i \in \text{Diff}_0(M, \mu)$ be diffeomorphisms satisfying each of the following conditions (we use here the notation of the proof of Theorem 1.2):

- the support of f_i and g_i is contained in the neighborhood of the union of the loops $\gamma_{i,j}$;
- the images $\Phi_i(f_i)$ and $\Phi_i(g_i)$ generate the free non-Abelian group in $\pi_1(M, m_i)/G_i$.

Such diffeomorphisms can be constructed in a similar way as g_i 's in the proof of Theorem 1.2.

Let $w \in \mathbf{F}_2$ be a reduced word and given two elements $f, g \in \text{Diff}(M, \mu)$ let $w(f, g)$ denote the induced diffeomorphism of M . Let

$$\Psi: \mathbf{F}_2 \times \dots \times \mathbf{F}_2 \rightarrow \text{Diff}(M, \mu, \sqcup_i B(m_i, r)) \subset \text{Diff}_0(M, \mu)$$

be defined by $\Psi(w_1, \dots, w_n) := w_1(f_1, g_1) \circ \dots \circ w_n(f_n, g_n)$. As before Ψ is a quasi-isometric embedding of a product of free groups on two generators into $\text{Diff}_0(M, \mu)$. Since \mathbf{F}_2 contains quasi-isometrically embedded a non-Abelian free group of an arbitrary finite rank [8] the proof is finished. \square

2.E. Proof of Theorem 1.4. The proof of the two dimensional case of Theorem 1.2 amounts to constructing a number of disjoint simple closed curves representing an undistorted element in the fundamental group of M . The present proof is analogous in the sense that we need to construct an embedding of the disjoint union of $2g + k - 2$ copies

of the figure eight into M such that each embedding induces a quasi-isometric embedding $\mathbf{F}_2 \rightarrow \pi_1(M, m_i)$ for $i = 1, \dots, 2g + k - 2$. We leave this straightforward construction as an exercise to the reader.

The rest of the proof is similar to the other proofs. That is, we construct relevant diffeomorphisms f_i, g_i and observe that the map

$$\Psi: \mathbf{F}_2 \times \cdots \times \mathbf{F}_2 \rightarrow \text{Diff}(\Sigma_{g,k}, \mu, \sqcup_i B(m_i, r)) \subset \text{Diff}_0(\Sigma_{g,k}, \mu)$$

defined by $\Psi(w_1, \dots, w_n) := w_1(f_1, g_1) \circ \cdots \circ w_n(f_n, g_n)$ is a quasi-isometric embedding. \square

Acknowledgments. We would like to thank Jim Howie and Mark Sapir for answering our questions. Example 1.7 is due to Jim Howie. This work has been done during the first author stay in Aberdeen. His visit was supported by the Coleman-Cohen foundation.

REFERENCES

- [1] Arnol'd V.: *Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits*. (French) Ann. Inst. Fourier (Grenoble) 16 1966 fasc. 1, 319–361.
- [2] Arnol'd V., Khesin B.: *Topological methods in hydrodynamics*, Applied Mathematical Sciences 125, (1998).
- [3] Benaïm M., Gambaudo J.-M.: *Metric properties of the group of area preserving diffeomorphisms*, Trans. Amer. Math. Soc. 353 no. 11, 2001, 4661–4672.
- [4] Bestvina M., Brady N.: *Morse theory and finiteness properties of groups*, Invent. Math. 129 (1997), 445–470.
- [5] Bridson M., Haefliger A.: *Metric spaces of non-positive curvature*, Springer-Verlag, New York, 1999.
- [6] Bridson M., Howie J., Miller C., Short H.: *Subgroups of direct products of limit groups*. Ann. of Math. (2) 170 (2009), no. 3, 1447–1467.
- [7] Crisp J., Wiest B.: *Quasi-isometrically embedded subgroups of braid and diffeomorphism groups*, Trans. Amer. Math. Soc. 359 2007, 5485–5503.
- [8] De la Harpe P.: *Topics in geometric group theory*, University of Chicago Press, 2000.
- [9] Dison W.: *Isoperimetric functions for subdirect products and Bestvina-Brady groups*, D.Phil. Thesis, Imperial College London, 2008. <http://www.maths.bris.ac.uk/~mawjd/thesis.pdf>
- [10] Eliashberg Y., Ratiu T.: *The diameter of the symplectomorphism group is infinite*, Invent. Math. 103 1991, 327–340.
- [11] Gottlieb D.: *A certain subgroup of the fundamental group*, Amer. J. Math. 87 1965, 840–856.
- [12] Kapovich M.: *RAAGs in Ham*, arXiv:1104.0348, 2011.
- [13] Kassel C., Turaev V.: *Braid groups*, Graduate Texts in Mathematics 247, Springer, 2008.
- [14] Khesin B., Wendt R.: *The geometry of infinite-dimensional groups*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 51. Springer-Verlag, Berlin, 2009.

- [15] Milnor J.: *A Note on Curvature and Fundamental Group*, J. Differential. Geometry 2, 1968.
- [16] Osin D.: *Small cancellations over relatively hyperbolic groups and embedding theorems*, Ann. of Math. (2) 172 (2010), no. 1, 1–39.
- [17] Roe J.: *Lectures in Coarse Geometry*, University Lecture Series Vol. 31, American Mathematical Society: Providence, Rhode Island, 2003.
- [18] Shnirelman A.: *Generalized fluid flows, their approximation and applications*. Geom. Funct. Anal. 4 (1994), no. 5, 586–620.

Department of Mathematics, Vanderbilt University, Nashville, TN 37240
E-mail address: michael.brandenburgsky@vanderbilt.edu

Department of Mathematics, Aberdeen University, Aberdeen, UK
E-mail address: kedra@abdn.ac.uk